

Horizontal transmission of a parasite is influenced by infected host phenotype and density

K. E. ROBERTS^{1*}† and W. O. H. HUGHES²

¹*School of Biology, University of Leeds, Leeds LS2 9JT, UK*

²*School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK*

(Received 29 January 2014; revised 30 May and 2 July 2014; accepted 6 July 2014)

SUMMARY

Transmission is a key determinant of parasite fitness, and understanding the dynamics of transmission is fundamental to the ecology and evolution of host–parasite interactions. Successful transmission is often reliant on contact between infected individuals and susceptible hosts. The social insects consist of aggregated groups of genetically similar hosts, making them particularly vulnerable to parasite transmission. Here we investigate how the ratio of infected to susceptible individuals impacts parasite transmission, using the honey bee, *Apis mellifera* and its microsporidian parasite *Nosema ceranae*. We used 2 types of infected hosts found simultaneously in colonies; sterile female workers and sexual males. We found a higher ratio of infected to susceptible individuals in groups resulted in a greater proportion of susceptibles becoming infected, but this effect was non-linear and interestingly, the ratio also affected the spore production of infected individuals. The transmission level was much greater in an experiment where the infected individuals were drones than in an experiment where they were workers, suggesting drones may act as intracolony ‘superspreaders’. Understanding the subtleties of transmission and how it is influenced by the phenotype of the infected/susceptible individuals is important for understanding pathogen transmission at population level, and for optimum targeting of parasite control strategies.

Key words: horizontal transmission, *Nosema ceranae*, honey bee, superspreader.

INTRODUCTION

Parasites impose a selection pressure on their hosts, influencing phenotypes and genotypes, and inducing changes at all levels of biological organization (Anderson and May, 1981; May and Nowak, 1994). Parasite fitness is inherently linked with the ability to transmit to new hosts, the mode and efficiency of which determine the success of any species or strain of parasite. The transmission dynamics of a parasite in a population depend on the interaction of host individuals, with the rate at which susceptible hosts become infected individuals being determined by the contact of susceptible hosts with infectious material and by the probability of the host becoming infected per contact (Begon *et al.* 2002).

For transmission to occur, a parasite must encounter a susceptible host. If the rate of transmission of new infections is proportional to the number of encounters between susceptible and infected hosts, then it can be modelled simply as βSI , where β is a constant representing the probability of contact and then probability of transmission, S is the number of susceptible individuals in a population and I is the

number of infected individuals in the population who may transmit the infection (Anderson and May, 1981). This is known as density-dependent transmission (Knell *et al.* 1998; McCallum *et al.* 2001; Begon *et al.* 2002). Both host number and density (hosts in an area) can also determine parasite fitness, which has led to the development of models that take into account changing transmission probability with changing host density (Anderson and May, 1981; De Jong, 1995; McCallum *et al.* 2001; Begon *et al.* 2002). Host density is at its most extreme in social animals, where there is spatial aggregation of potential hosts, most notably in the social insects where within a colony there is also temporal overlapping of generations of genetically similar hosts (Schmid-Hempel, 1998). Consequently, sociality is often predicted to be associated with increased disease transmission (Freeland, 1976; Shykoff and Schmid-Hempel, 1991; Johnson *et al.* 2011), although this effect may be mitigated by a decrease in intergroup transmission and highly evolved group-level defences (Hughes *et al.* 2002; Traniello *et al.* 2002; Wilson *et al.* 2003; Boomsma *et al.* 2005).

In social insects, within-group transmission can be at low levels with the transmission coefficient decreasing with increasing host density (Rosengaus *et al.* 2000; Hughes *et al.* 2002; Traniello *et al.* 2002). It has been suggested that the social organization within a social insect colony could have evolved to reduce the spread of pathogens, an idea referred to as

* Corresponding author: School of Biology, University of Leeds, Leeds LS2 9JT, UK. E-mail: k.roberts@exeter.ac.uk

† Current address: College of Life and Environmental Sciences, University of Exeter, Cornwall Campus, Penryn TR10 9EZ, UK.

organizational immunity (Naug and Camazine, 2002; Naug and Smith, 2007). However, social insect colonies, as with groups of any animal, include multiple phenotypes (castes and ages) and genotypes, which differ in the resistance to parasites (e.g. Baer and Schmid-Hempel, 2003; Baer *et al.* 2005; Poulsen *et al.* 2006; Ruiz-González and Brown, 2006; Armitage and Boomsma, 2010; Hughes *et al.* 2010; Smart and Sheppard, 2012; Roberts and Hughes, 2014), something that may have profound effects on the within-group dynamics of parasite transmission. Social insects therefore make ideal models for studying parasite transmission, because their lifestyle of living in dense, highly social groups of related individuals with varying phenotypes and genotypes provides an opportunity to test a multitude of social transmission processes (Schmid-Hempel, 1998).

Here we investigate the dynamics of horizontal transmission using the honey bee *Apis mellifera* and its microsporidian parasite *Nosema ceranae* as the model system. *Nosema ceranae* is horizontally transmitted between adult bees through ingestion of infective spores which germinate and infect gut epithelial cells, before proliferating and producing more infective spores which are excreted by the host in the feces, with subsequent transmission to susceptible individuals being either fecal–oral or oral–oral (Fries, 2010; Smith, 2012). The dynamics of *N. ceranae* transmission are not fully understood, due to its relatively recent and apparently rapid global colonization of *A. mellifera* (Higes *et al.* 2006; Klee *et al.* 2007). The parasite has, however, been found to infect all castes of adult honey bee: the sterile female workers (Higes *et al.* 2007; Fries, 2010), the sexual male drones and sexual female queens (Traver and Fell, 2011a, b; Botías *et al.* 2012). We investigate how transmission rate relates to the ratio of infected to susceptible individuals in a group, by comparing parasite transmission in groups with infected:susceptible (I:S) ratios of 5:5, 4:6, 3:7, 2:8, 1:9, as well as 0:10 control groups. We do this in separate experiments for groups where the infected individuals were either workers or drones, to examine whether the phenotype of infected individuals may alter the transmission relationship. The null hypothesis if the transmission parameter β was constant would be that the number of susceptible individuals which become infected is proportional to IS , and differ respectively by the ratio 25:24:21:16:9 for the I:S ratios we test here.

MATERIALS AND METHODS

Experimental hosts and parasite preparation

Bees were obtained from managed *A. mellifera* colonies of similar age and size. Bees were obtained from 10 colonies for the experiment in which workers were the experimentally infected individuals and from 9 colonies for the drone experiment, with 4 of the

colonies providing bees for both (see Supplementary material). All colonies were confirmed to be free from *Nosema* infection by screening 30 returning foragers for the presence of *Nosema* by light microscopy and conventional polymerase chain reaction (PCR) (Chen *et al.* 2008; see below for details; no control bees developed *Nosema* infections during the experiment, confirming that this methodology was robust for identifying the general infection status of a colony). To obtain experimental bees of the same age, frames of brood from the colonies were allowed to eclose in incubators at 34 °C, 60% RH, 12:12 light:dark cycle, with newly hatched bees taken from the frame and placed into plastic pots (Ambican 16oz Fabrikal Delipot, 115 × 95 × 75 mm) with sterile filter paper on the base of the pots and 50% sterile sucrose solution provided *ad libitum*. Bees were kept in incubators under the same conditions, in the same pots throughout the whole experiment. Parasite spores were isolated from 100 *A. mellifera* foragers collected from 5 naturally infected colonies. The ventriculi of the bees was homogenized in 1.5 mL microcentrifuge tubes containing 500 μ L molecular grade water using a plastic pestle, and then examined by light microscopy to confirm the presence of *Nosema* spores. Conventional PCR was then used to identify the species of *Nosema* (see below). Spore suspensions from bees that were infected by *N. ceranae* only, through molecular screening, were purified with Percoll centrifugation using a modification of Huang *et al.* (2007). The concentration of spores was quantified using FastRead (Immune Systems) disposable haemocytometers. Spore viability was checked using 10% Trypan blue dye staining (Martin-Hernandez *et al.* 2009), to allow spore doses to be adjusted to account for non-viable spores. Spores were then diluted in a sterile 50% sucrose solution to make up an infective dose of 2500 viable spores per μ L. Fresh spore suspensions were prepared from the same colonies and used for the infections of the drones and workers, respectively, within an interval of 10 days.

Experimental infection

The experiment compared I:S individuals of 5:5, 4:6, 3:7, 2:8, 1:9 or 0:10 (the control groups), with infected individuals being workers in 1 experiment and drones in a second experiment and the susceptible individuals being workers in both (see Fig. S1 for experimental design). To generate infected individuals, 6-day-old bees were individually hand-fed with 10 000 viable *N. ceranae* spores in a 4 μ L sterile sucrose solution using a micropipette. Bees were held in 0.5 μ L microcentrifuge tubes with the ends cut off during this process, to reduce their stress levels by minimizing handling and removing the need for anaesthetic. Bees were starved for 1 h before infection to ensure they would fully ingest their dose. Any bees

that did not fully ingest their dose were discarded and replaced by another individual. All treated bees were found to be infected by *N. ceranae* by PCR screening at the end of the experiment and so are described as infected individuals hereafter. Two sets of controls were used throughout the experiments: (a) 5 sham-treated individuals that were fed 4 μL of the sterile sucrose solution and (b) unmanipulated bees. Sterile sucrose solution was used rather than a *Nosema*-negative Percoll washed bee homogenate; previous optimization of methods showed no difference in survival between bees fed *Nosema*-negative Percoll washed extract and sucrose solution (Huang *et al.* 2007; Chen *et al.* 2009; Cornman *et al.* 2009; Martin-Hernandez *et al.* 2009). Bees were held individually for an hour post-inoculation to minimize bee-to-bee (trophallactic) sharing of their dose, before being placed in groups of like-treated nestmates for 24 h to allow the inoculum to be digested (Crailsheim, 1988). After this 24 h period, infected bees were set up in plastic pots with untreated, susceptible bees from their respective colonies to create groups of 10 bees with I:S ratios of 5:5, 4:6, 3:7, 2:8, 1:9 or 0:10 (the control groups). There was 1 group per colony for each I:S ratio, giving 10 replicate groups (i.e. from 10 different colonies) of each ratio when the experimentally infected individuals were workers and 9 replicate groups (i.e. from 9 different colonies) of each ratio when they were drones (Fig. S1). Both infected and susceptible bees were marked with different colour paint dots to allow them to be identified subsequently. The survival of the bees was then monitored daily for 12 days, to allow sufficient time for any transmitted infections to develop whilst avoiding excessive bee mortality. Bees that died during this period, as well as all bees surviving at the end of the experiment, were stored in 90% ethanol at -20°C for subsequent quantification of parasite infection.

Parasite detection

To determine the infection status of individuals, the ventriculi of bees were individually homogenized in 500 μL of molecular grade water using a plastic pestle. The numbers of *Nosema* spores were then quantified using FastRead counting chambers, and the remainder of the homogenate used for species-specific PCR analysis of *Nosema*. DNA was extracted by incubating 75 μL of the ventriculus homogenate for 12 h with Proteinase K (Promega) ($5\ \mu\text{L}\ \text{mL}^{-1}$) at 56°C , and then boiling with 75 μL of 5% Chelex 100 (Biorad) suspended in 10 μM Tris Buffer. After centrifugation, the supernatant was stored at -20°C for molecular analysis. Molecular detection was carried out using *N. ceranae* and *Nosema apis* specific primers from Chen *et al.* (2008), and the quality of DNA extractions was checked using the *LCO-Hym/HCOout* primers for the CO1 host control gene (Folmer *et al.* 1994; Prendini *et al.* 2005). To ensure accuracy

of *Nosema* detection, a 20% subsample of positive and negative samples from each colony were double checked with additional primers targeting species-specific regions of the *RPB1* gene (Gisder and Genersch, 2013), which gave from the same results as those found using the Chen *et al.* (2008) primers. Reactions were carried out in 96-well microtitre plates in a VertiPCR thermal cycler (ABI), with primer-specific mixes containing 2.5 mM MgCl_2 and 2.5 mM dNTPs, 0.2 μM of each primer, 0.8 U Taq (PROMEGA) and 50–100 ng of DNA in a 10 μL reaction. PCR thermo-cycling profiles for all reactions consisted of an initial denaturing step of 94°C for 2 min, followed by 35 cycles of 94°C for 30 s, annealing temperature for 45 s and 72°C for 2 min, with a final elongation step of 72°C for 7 min. The annealing temperatures were 50°C for the host control gene, 61°C for *N. ceranae* and 63°C for *N. apis*. PCR products were visualized under UV using 1% agarose gels stained with ethidium bromide and compared with a 100 base pair, size ladder. All plates were run with both triplicate negative and positive controls, and all samples were run twice, which in all cases confirmed that negatives were true negatives.

Statistical analysis

All analysis was carried out in R 2.14.2 (R Development Core Team, 2012). As the workers and drones were tested in separate experiments, the 2 datasets were analysed separately. Survival analysis was carried out using the *survreg* function in the Survival package using default Weibull errors to account for non-constant errors. Frailty models were used in order to include colony as a random factor, with day of death as the response variable and treatment, age and their interaction as the variables. Hazard ratios were extracted from survival models of the hazard of death of infected bees *vs* the untreated control bees. As there were no differences in survival between the 2 sets of controls (sham-treated bees and unmanipulated bees) they were therefore pooled for calculating the hazard ratios. Transmission potential, as estimated by spore counts, and infection status determined by PCR were both analysed using mixed effects models implemented using the *lme4* package (Bates and Maechler, 2010). Either the spore count or the infection status was fitted as the response variable, with the ratio of I:S individuals as the discrete independent variable and colony as a random factor. Spore counts were modelled with a *glmer* to allow data to be fitted with a Poisson error structure and included the day of death of each bee to control for any effect of duration of infection on infection intensity. Infection status was modelled as a binomial response (infected or uninfected). Significance was based on model comparison using likelihood ratio tests comparing models of fixed effects to null models

with only random effects. Where appropriate we accepted models where the model with fixed effects differed significantly from the null model, and therefore report the total degrees of freedom for the minimal model, as well as the P -value associated with model comparison. Models containing only 1 fixed variable did not require model selection. All models were checked for normality of errors, and homogeneity of variance, through plots of residuals against fitted values and quantile–quantile plots, and no data or models violated the assumptions of analysis. In addition, for the 4 colonies that were used as source colonies for both the drone and worker experiments (see Supplementary material S1), the survival, proportion infected and spore production of the susceptible and infected individuals were analysed together, as described for the separate experiments. This analysis found similar differences between workers and drones to those suggested by comparison of the overall experiments, suggesting that these differences were not simply due to difference in the colonies from which workers and drones originated (see Supplementary material S2). Transmission terms were calculated for each cage of infected to susceptible ratio, and averaged across colonies for each worker and drones treatment, using the simple, mass action model, βSI (Knell *et al.* 1996; Begon *et al.* 2002; McCallum *et al.* 2002). As we have experimentally controlled population number and area, here S is the number of susceptible hosts, I is the number of infected hosts and β is the transmission coefficient. Calculations of β were also carried out using the number of spores produced by the experimentally infected individuals. Correlations between the transmission coefficient and the I:S ratios were carried out using linear regressions.

RESULTS

Spore production of experimentally infected individuals

All experimentally infected bees were found to have become infected with *N. ceranae*, being positive by both PCR screening and visual detection of spores in their ventriculi. None of the control bees were found by light microscopy or PCR to have any *Nosema* infections. The numbers of spores that the experimentally infected workers and drones produced was highly variable; however, the spore production of workers was significantly affected by the ratio of I:S bees in a group, despite all the infected bees having received the same initial dose of parasite ($\chi^2_{4,7} = 12.42$, $P = 0.014$, with control groups excluded; Fig. 1a). Infected workers in groups with between 2 and 5 infected individuals had similar numbers of spores, whereas those in the groups with only a single infected individual produced a much higher number of spores (Fig. 1a). The number of experimentally infected drones in a group was also found to

significantly affect the number of spores that they produced ($\chi^2_{4,7} = 14.53$, $P = 0.005$, Fig. 1b). Experimentally infected drones had more variation in the average number of spores they produced than did infected workers (Fig. 1), and did not have the high spore production in the groups with a single infected individual that was shown by groups in which the infected individual was a worker.

Infection of susceptible bees

For the susceptible individuals housed with either experimentally infected workers or drones, there was a significant effect of the ratio of I:S bees on the proportion of the susceptible individuals that became infected (when the experimentally infected individuals were workers: $\chi^2_{4,7} = 45.7$, $P < 0.001$, Fig. 2a; when the experimentally infected were drones: $\chi^2_{4,7} = 99.8$, $P < 0.001$, Fig. 2b). The effect of a decrease in the I:S ratio on the numbers of susceptible individuals becoming infected was substantially less than expected if the transmission parameter β was constant (Fig. 2). Susceptible bees were more likely to become infected when the proportion of experimentally infected individuals in their group was greater (Fig. 2). However, there was a dramatic difference between the worker and drone experiments, with many more susceptible individuals becoming infected when they were kept with infected drones than when kept with infected workers (Fig. 2a and b). All susceptible bees in the control groups were found by PCR to be negative for *N. ceranae* and no spores were found in them when inspected by light microscopy. As with the experimentally infected individuals, the spore production of the susceptible individuals was highly variable. However, the proportion of experimentally infected individuals, either worker or drones, also affected the spore production of susceptible individuals ($\chi^2_{5,11} = 9.82$, $P = 0.028$, Fig. 1c, and $\chi^2_{2,6} = 11.29$, $P = 0.003$, Fig. 1d, respectively). Susceptible bees that were housed with 2–4 infected workers produced around 6×10^5 spores, whereas those with 1 or 5 infected workers had 6×10^6 spores. There was large variation in the spore numbers produced in some treatments. Those housed with 2–5 infected drones had higher numbers of spores than those housed with infected workers, whereas those with a single infected drone had relatively few spores. The variation is partially due to some individuals having very few spores present, or in the case of the 3:7 and 4:6 ratios, is due to some bees (4 and 2 individuals, respectively) having no spores counted despite PCR screening indicating that they were infected.

Survival

Survival was significantly different between those experimentally infected with *N. ceranae* and the




Fig. 1. Mean (\pm S.E.) number of spores of the *N. ceranae* parasite in the ventriculi of experimentally infected (a) workers or (b) drones which had been fed a dose of 1×10^4 *N. ceranae* spores, and the susceptible workers kept with the experimentally infected (c) workers or (d) drones for 12 days. Bees were kept in groups with a 5:5, 4:6, 3:7, 2:8 or 1:9 ratio of infected to susceptible individuals. Only bees confirmed by PCR to be infected by *N. ceranae* are included in the spore counts (all experimentally infected bees and see Fig. 2 for proportion of the susceptible workers which became infected).

control groups for both the workers and drones ($\chi^2_1 = 20.5$, $P < 0.001$ and $\chi^2_1 = 12.9$, $P < 0.001$, respectively), with infection of *Nosema* causing higher mortality. Survival of the experimentally infected bees differed significantly depending on the ratio of susceptible to infected bees, both when the infected bees were workers and drones ($\chi^2_5 = 16.6$, $P = 0.005$ and $\chi^2_{11,7} = 65.8$, $P < 0.001$, respectively; Fig. 3, Supplementary Figs. S2 and S3). Experimentally infected bees in lower ratios of infected to susceptible individuals appeared to survive better (Fig. 3). The survival of susceptible individuals in groups with experimentally infected workers was not found to significantly influence the proportion of infected individuals ($\chi^2_5 = 0.24$, $P = 1.0$, Fig. 3c), with all groups having a similarly low hazard from *N. ceranae* exposure and showing very marginally better survival than control bees (hence the hazard ratios of slightly < 1). However, the survival of susceptible individuals kept with the experimentally infected drones was markedly different, being significantly influenced by ratio of susceptible to infected individuals

($\chi^2_{11,9} = 69.2$, $P < 0.001$, Fig. 3d), with the survival of susceptible individuals kept with infected drones being substantially worse overall than when kept with infected workers.

Transmission coefficients

When calculated using experimentally infected individuals as *I*, the transmission coefficients had a negatively association with the I:S ratio when experimentally infected individuals were workers ($F_{1,46} = 8.14$, $P = 0.06$, $R^2 = 0.64$), but positively correlated when they were drones ($F_{1,46} = 11.4$, $P = 0.004$, $R^2 = 0.72$, Fig. 4a). Transmission coefficients were substantially lower when the numbers of parasite spores produced by experimentally infected individuals were used as *I*, and in this case were negatively correlated with the I:S ratio with the exception of the 1:9 ratio in the drone experiment when the transmission coefficient was much greater (worker experiment: $F_{1,46} = 40.1$, $P = 0.007$, $R^2 = 0.90$; drone experiment: $F_{1,46} = 1.5$, $P = 0.3$,




Fig. 2. Mean (\pm s.e.) proportion of susceptible worker bees who became infected with the *N. ceranae* parasite after 12 days housed with experimentally infected (a) workers or (b) drones, and kept in groups with either a 5:5, 4:6, 3:7, 2:8 or 1:9 ratio of infected to susceptible individuals. Dashed line represents the expected numbers of new infections that would occur if the transmission coefficient was constant, shown here for the value calculated for the 5:5 ratio.

$R^2 = 0.90$, Fig. 4b). The transmission coefficients were in general higher when experimentally infected individuals were drones than when they were workers in both cases.

DISCUSSION

The ratio of infected to susceptible individuals in a group was positively related to the intensity and prevalence of infections in susceptible individuals, but the relationship was much weaker than expected if the transmission parameter β was constant. The effect of the ratio of susceptible to infected individuals appeared to differ depending on the infected phenotype, with more transmission occurring in the experiment in which the infected individuals were drones than when they were workers. Interestingly, the spore production of the experimentally infected bees, whilst highly variable, was not constant across the treatment groups, despite them all having received the same initial parasite dose, suggesting that either the ratio of I:S individuals in a group or the density of infectious hosts affects not only the dynamics of between-host transmission, but also the within-host infection. Although it is possible that some secondary infections may have contributed to the results, these are likely to have been limited because spore production of *N. ceranae* is low until at least 6 days post-infection (Paxton *et al.* 2007; Forsgren and Fries, 2010; Fries, 2010).

We found that the ratio of I:S individuals influenced the subsequent infection rate of the susceptible individuals and the numbers of susceptible

individuals that became infected at the lower I:S ratios were less than expected if the transmission parameter β was a constant. Horizontal transmission of parasites is often dependent on contact between infected and susceptible individuals and is therefore influenced by the ratio of these in a population (Anderson and May, 1981, 1982; McCallum *et al.* 2001, 2002). When group size is constant, higher ratios of I:S individuals are predicted to result in higher proportions of susceptible individuals becoming infected (Knell *et al.* 1996; Ryder *et al.* 2007), as found to be the case here. This result indicates that a density-dependent form of transmission may occur in *N. ceranae* (McCallum *et al.* 2001, 2002; Begon *et al.* 2002). Interestingly, the effect of the I:S ratio on the probability of susceptible individuals becoming infected with either workers or drones did not match that predicted if the transmission parameter was constant. In some host–parasite systems the relationship asymptotes (Fenton *et al.* 2002), but in this case there was a substantial increase in the probability between I:S ratios of 4:6 and 5:5. Potentially this may be due to the increasing infection pressure on the susceptible individuals in these cohorts, as the infection prevalence increases so too may the concentration of parasite propagules in the environment (Matthews *et al.* 2009). Individual bees may be able to handle low I:S ratios, avoiding infection by employing individual and social immunity; however, once the ratio becomes more extreme these defences may be overcome.

The infected to susceptible ratio also influenced the intensity of the infection seen in the susceptible




Fig. 3. Survival graphs of individuals over a 12-day period after exposure to the *N. ceranae* parasite for (a) experimentally infected worker bees or (b) experimentally infected drones, kept for 12 days with susceptible workers (c) those kept with infected workers and (d) kept with infected drones at either a 5:5, 4:6, 3:7, 2:8 or 1:9 ratio of infected to susceptible individuals (control: solid black line; 1:9 grey small dash line; 2:8 grey large dash; 3:7 black small dash; 4:6 grey solid line and 5:5 black small dash line). Corresponding hazard ratio graphs for each group are included as inset graphs, calculated as the hazard of exposure in comparison to the control groups. Note the difference in y-axis scale between the graphs.

individuals once they became infected. Higher infection levels in susceptible individuals may occur due to multiple infection events, as susceptible individuals in the groups with higher ratios of infected individuals may encounter a larger number of transmission events, effectively receiving a higher overall dose and therefore infection intensity. Interestingly the ratio of infected to susceptible individuals in the group also influenced the intensity of infection seen in the experimentally infected bees, despite them all receiving the same initial dose. In workers, spore production was much higher at an I:S ratio of 1:9 than at higher ratios, whereas in drones the effect was inconsistent, which may in part be due to the greater variation in spore production shown by drones at most of the I:S ratios, with some individuals producing up to 5 times less spores. The effect, however, was significant in both phenotypes, in spite of the variation in the data, so it appears that the I:S ratio does alter the spore production of infected individuals, but not necessarily in a straightforward way.

The higher spore production rates may also be due to behavioural differences in the bees. Any individuals that are detected as being infected may be subject to more aggressive treatment or behavioural stressors, or receive less care and social interactions from their nest mates (Moore, 1995; Cremer *et al.* 2007; Cotter and Kilner, 2010; Rueppell *et al.* 2010). The cuticular hydrocarbon profile of infected individuals has been known to change following an immune challenge and therefore allow infection status to be determinable by nest mates (Richard *et al.* 2008, 2012). The infected individuals may potentially be segregated behaviourally from the other individuals to some extent to minimize parasite transmission, thus influencing the transmission dynamics of the parasite in a context-dependent way. It may be that the greater variation and inconsistency in the spore production of drones was due to them relying on nestmate workers for food and being treated less consistently by them than was the case for other workers. *Nosema* infection has also been linked to an




Fig. 4. Transmission parameter of infected workers (a, b) and drones (c, d) housed with susceptible workers in ratios of 5:5, 4:6, 3:7, 2:8 or 1:9 susceptible to infected individuals (a) workers and (c) drones calculated using the number of individuals as the infectious unit and calculated using the spore counts as the infectious units (b) worker and (d) drone, error bars representing the mean standard error of the transmission parameter for each pot of bees.

increase in hunger and behavioural changes in infected individuals, so it is also possible that infected bees may try to engage in more trophallaxis than susceptible bees in order to counter-act the energetic demand of infection (Mayack and Naug, 2009; Naug and Gibbs, 2009).

The transmission rate appeared to be greater in the experiment in which drones were the infected individuals than in the experiment in which the infected individuals were workers. As the drones and workers were tested in separate experiments, it cannot be certain that this reflects a genuine difference in transmission rate between phenotypes, but this seems feasible. The drone and worker experiments were carried out under identical conditions with only a 10-day period between them. While different suspensions of the parasite were used in order to control for viability, these were obtained from the same host colonies and it is unlikely that the genetic composition or infection dynamics of the parasite changed in such a short period. The difference also seems unlikely to be due to differences in the genetic composition of the hosts, because the same effect was

present in the subset of colonies that were used in both drone and worker experiments, as in the overall samples (Supplementary information). It is nevertheless possible that some other unknown factor may have affected the results, such as parasite life cycle differences between the host types, and further work is needed to confirm other possible causes. It, however, seems likely that there may be a phenotypic difference in transmission rate in honey bees. Male honey bees have been shown to have increased parasitism rates by the mite *Varroa destructor*, a known vector of viruses such as Deformed Wing Virus (Fuchs, 1990; Boot *et al.* 1995; de Miranda and Fries, 2008). It is therefore possible that other parasites, as well as *Nosema*, may have been stressing the drones and thereby in part causing the greater infectivity of drones with *Nosema*. Sexual dimorphism in transmission rate has previously been attributed to high parasite load or elevated propagule production rate (Perkins *et al.* 2003; Galvani and May, 2005; Gear *et al.* 2012), and to males investing less in immunity (Bateman, 1948; Rolff, 2002; Nunn *et al.* 2009). In the case of honey bees, males may also be more

susceptible because they are haploid (O'Donnell and Beshers, 2004) or receive more food by trophallaxis (Higes *et al.* 2009; Feigenbaum and Naug, 2010; Traver and Fell, 2011a, b; Smith, 2012), and it is also possible that the cell cycle of *N. ceranae* may be shorter in males than workers. Regardless of the mechanism responsible, if drones do transmit the parasite far more than workers, this may make drones potential 'superspreaders' of infection within a colony (Woolhouse *et al.* 1997; Lloyd-Smith *et al.* 2005). Pathogen load and shedding rates of parasites are rarely measured in superspreading events (Lloyd-Smith *et al.* 2005), so the heterogeneity in these components that is suggested by our results may potentially be important. Identifying these influential hosts in transmission is beneficial to understanding parasite spread and control, as treatment strategies would be most effective if targeting these individuals (Woolhouse *et al.* 1997; Lloyd-Smith *et al.* 2005; Hawley and Altizer, 2011).

Transmission of parasites is dependent on contact between infected and susceptible individuals, and therefore the ratio of these in a population. From our data we calculated the transmission coefficient β from a mass action model, with β describing the rate at which susceptible individuals are converted to infected hosts (due to the contact rate and probability of infection per contact; Begon *et al.* 2002). This revealed that the transmission parameter was positively correlated with the I:S ratio, which contrasts with the negative relationship seen in other studies (e.g. Knell *et al.* 1998). There are at least 3 non-mutually exclusive hypotheses that could explain this. First, it may be that *Nosema* infections show an allee effect, with a minimum number of spores needing to be ingested for infections to be likely to be successful, as is the case for other parasites (Hughes *et al.* 2004). Second, bees may isolate or avoid interacting with infected individuals which would reduce transmission and such social exclusion may be more effective when the I:S ratio is low. Third, lower frequencies of infected individuals may lead to greater spatial heterogeneity in the deposition of spores and consequently reduced likelihood of susceptible individuals encountering spores when movement in the environment is non-random, although this effect is unlikely in the small containers used in this experiment. The transmission parameter also appeared to be higher when the infected individuals were drones rather than workers, and that the relationship with I:S ratio may also differ between phenotypes. The mechanism of drone transmission is likely to be through a mixture of fecal-oral and oral-oral transmission (Higes *et al.* 2008; Smith, 2012). Although we did not carry out quantitative behavioural observations, feces was seen on the filter paper floors of the cages, confirming that defecation took place and thus the potential for the ingestion of spores from feces during grooming or hygienic

behaviours. Huang and Solter (2013) also commented that they found *Nosema* spores present on the inside of cages, and that spores found in the mouth of bees were likely due to cleaning and grooming behaviours. We observed some drone-to-drone exchange of food during the experimentally set up, when drones were held together for 24 h before being placed with the naive susceptible individuals. Although this behaviour may be rare it suggests that drones will initiate and take part in feeding others, with the potential for parasite transmission. Phenotype-dependent transmission coefficients have been demonstrated previously in *Plodia interpunctella* infected with *B. thuringiensis*, in which 5th instar larvae have a higher transmission term than infected 4th instars (Knell *et al.* 1996). This highlights the fact that in a given host-parasite system, transmission dynamics may differ depending on not just the host and/or parasite genetics. The transmission terms do not appear to have a fully linear relationship, which shows, unsurprisingly, that the simple model used here is not sufficient to fully explain the dynamics of this system. Interestingly, the transmission coefficients were substantially lower when calculated using the numbers of parasites spores as I , rather than the number of infected hosts. The very low transmission coefficients for spores indicate the relative inefficiency of transmission in this parasite.

The results emphasize the importance of recognizing the heterogeneity in host-parasite transmission dynamics. The ratio of infected to susceptible individuals can influence not only the probability of a susceptible becoming infected, but also the dynamics of the within-host infection and resulting production of parasite propagules by an infected host, with both also potentially being affected by host phenotype. Understanding the dynamics of parasite transmission, and how this can change with host phenotype, is likely to be fundamental to our comprehension of parasite virulence, as well as for disentangling reciprocal cycles in population sizes.

SUPPLEMENTARY MATERIAL

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S0031182014001243>.

ACKNOWLEDGEMENTS

We thank Bill Cadmore, Sophie Evison and Crystal Frost for technical assistance, Peter Graystock, Jon Ryder and Ben Jones for helpful discussion, and Sophie Evison and 3 anonymous reviewers for valuable comments which improved the manuscript.

FINANCIAL SUPPORT

This work was supported by a Biotechnology and Biological Sciences Research Council studentship to KER.

REFERENCES

- Anderson, R. M. and May, R. M. (1981). The population dynamics of microparasites and their invertebrate hosts. *Philosophical Transactions of the Royal Society of London Biological Sciences* **291**, 451–524. doi: 10.1098/rstb.1981.0005.
- Anderson, R. M. and May, R. M. (1982). Coevolution of hosts and parasites. *Parasitology* **85**, 411–426.
- Armitage, S. A. O. and Boomsma, J. J. (2010). The effects of age and social interactions on innate immunity in a leaf-cutting ant. *Journal of Insect Physiology* **56**, 780–787. doi: 10.1016/j.jinsphys.2010.01.009.
- Baer, B. and Schmid-Hempel, P. (2003). Bumblebee workers from different sire groups vary in susceptibility to parasite infection. *Ecology Letters* **6**, 106–110. doi: 10.1046/j.1461-0248.2003.00411.x.
- Baer, B., Krug, A., Boomsma, J. J. and Hughes, W. O. H. (2005). Examination of the immune responses of males and workers of the leaf-cutting ant *Acromyrmex echinator* and the effect of infection. *Insectes Sociaux* **52**, 298–303. doi: 10.1007/s00040-005-0809-x.
- Bateman, A. (1948). Intra sexual selection in *Drosophila*. *Heredity* **2**, 349–368. doi: 10.1038/hdy.1948.21.
- Bates, D. and Maechler, M. (2010). lme4: linear mixed-effects models using Eigen and Eigen. *R package version 0.999375–33*.
- Begon, M., Bennett, M., Bowers, R. G., French, N. P., Hazel, S. M. and Turner, J. (2002). A clarification of transmission terms in host-microparasite models: numbers, densities and areas. *Epidemiology and Infection* **129**, 147–153. doi: 10.1017/S0950268802007148.
- Boomsma, J. J., Schmid-Hempel, P. and Hughes, W. O. H. (2005). Life histories and parasite pressure across the major groups of social insects. In *Insect Evolutionary Ecology* (ed. Fellowes, M. D. E., Holloway, G. J. and Rolff, J.), pp. 139–176. CABI Publishing, London.
- Boot, W. J., Schoenmaker, J., Calis, J. N. M. and Beetsma, J. (1995). Invasion of *Varroa jacobsoni* into drone brood cells of the honey bee, *Apis mellifera*. *Apidologie* **26**, 109–118. doi: 10.1051/apido:19950204.
- Botías, C., Martín-Hernández, R., Dias, J., García-Palencia, P., Matabuena, M., Juarranz, A., Barrios, L., Meana, A., Nanetti, A. and Higes, M. (2012). The effect of induced queen replacement on *Nosema* spp. infection in honey bee (*Apis mellifera iberiensis*) colonies. *Environmental Microbiology* **14**, 845–859. doi: 10.1111/j.1462-2920.2011.02647.x.
- Chen, Y. P., Evans, J. D., Smith, I. B. and Pettis, J. S. (2008). *Nosema ceranae* is a long-present and wide-spread microsporidian infection of the European honey bee (*Apis mellifera*) in the United States. *Journal of Invertebrate Pathology* **97**, 186–188. doi: 10.1016/j.jip.2007.07.010.
- Chen, Y. P., Evans, J. D., Murphy, C., Gutell, R., Zuker, M., Gundensen-Rindal, D. and Pettis, J. S. S. (2009). Morphological, molecular, and phylogenetic characterization of *Nosema ceranae*, a microsporidian parasite isolated from the European honey bee, *Apis mellifera*. *The Journal of Eukaryotic Microbiology* **56**, 142–147. doi: 10.1111/j.1550-7408.2008.00374.x.
- Cornman, R. S., Chen, Y., Schatz, M. C., Street, C., Zhao, Y., Desany, B., Egholm, M., Hutchison, S., Pettis, J. S., Lipkin, W. I. and Evans, J. D. (2009). Genomic analyses of the microsporidian *Nosema ceranae*, an emergent pathogen of honey bees. *PLoS Pathogens* **5**, e1000466. doi: 10.1371/journal.ppat.1000466.
- Cotter, S. C. and Kilner, R. M. (2010). Personal immunity versus social immunity. *Behavioral Ecology* **21**, 663–668. doi: 10.1093/beheco/arq070.
- Crailsheim, K. (1988). Regulation of food passage in the intestine of the honeybee. *Journal of Insect Physiology* **34**, 85–90. doi: 10.1016/0022-1910(88)90158-8.
- Cremer, S., Armitage, S. A. O. and Schmid-Hempel, P. (2007). Social immunity. *Current Biology* **17**, R693. doi: 10.1016/j.cub.2007.06.008.
- De Jong, M. C. M. (1995). Mathematical modelling in veterinary epidemiology: why model building is important. *Preventive Veterinary Medicine* **25**, 183–193. doi: 10.1016/0167-5877(95)00538-2.
- De Miranda, J. R. and Fries, I. (2008). Venereal and vertical transmission of deformed wing virus in honeybees (*Apis mellifera* L.). *Journal of Invertebrate Pathology* **98**, 184–189. doi: 10.1016/j.jip.2008.02.004.
- Feigenbaum, C. and Naug, D. (2010). The influence of social hunger on food distribution and its implications for disease transmission in a honeybee colony. *Insectes Sociaux* **57**, 217–222. doi: 10.1007/s00040-010-0073-6.
- Fenton, A., Fairbairn, J. P., Norman, R. and Hudson, P. J. (2002). Parasite transmission: reconciling theory and reality. *Journal of Animal Ecology* **71**, 893–905.
- Folmer, O., Black, M., Hoeh, W., Lutz, R. and Vrijenhoek, R. (1994). DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* **3**, 294–299.
- Forsgren, E. and Fries, I. (2010). Comparative virulence of *Nosema ceranae* and *Nosema apis* in individual European honey bees. *Veterinary Parasitology* **170**, 212–217. doi: 10.1016/j.vetpar.2010.02.010.
- Freeland, W. J. (1976). Pathogens and the evolution of primate sociality. *Biotropica* **8**, 12–24.
- Fries, I. (2010). *Nosema ceranae* in European honey bees (*Apis mellifera*). *Journal of Invertebrate Pathology* **103**(Suppl.), S73–S79. doi: 10.1016/j.jip.2009.06.017.
- Fuchs, S. (1990). Preference for drone brood cells by *Varroa jacobsoni* Oud in colonies of *Apis mellifera carnica*. *Apidologie* **21**, 193–199. doi: 10.1051/apido:19900304.
- Galvani, A. P. and May, R. M. (2005). NEWS and VIEWS dimensions of superspreading. *Nature* **438**, 293–295. doi: 10.1038/438293a.
- Gisder, S. and Genersch, E. (2013). Molecular differentiation of *Nosema apis* and *Nosema ceranae* based on species-specific sequence differences in a protein coding gene. *Journal of Invertebrate Pathology* **113**, 1–6. doi: 10.1016/j.jip.2013.01.004.
- Grear, D. A., Luong, L. T. and Hudson, P. J. (2012). Sex-biased transmission of a complex life-cycle parasite: why males matter. *Oikos* **121**, 1446–1453. doi: 10.1111/j.1600-0706.2012.20358.x.
- Hawley, D. M. and Altizer, S. M. (2011). Disease ecology meets ecological immunology: understanding the links between organismal immunity and infection dynamics in natural populations. *Functional Ecology* **25**, 48–60. doi: 10.1111/j.1365-2435.2010.01753.x.
- Higes, M., Martín-Hernández, R., Aránzazu, M., Martín, R. and Meana, A. (2006). *Nosema ceranae*, a new microsporidian parasite in honeybees in Europe. *Journal of Invertebrate Pathology* **92**, 93–95. doi: 10.1016/j.jip.2006.02.005.
- Higes, M., García-Palencia, P., Martín-Hernández, R. and Meana, A. (2007). Experimental infection of *Apis mellifera* honeybees with *Nosema ceranae* (Microsporidia). *Journal of Invertebrate Pathology* **94**, 211.
- Higes, M., Martín-Hernández, R., Botías, C., Bailón, E. G., González-Porto, A. V., Barrios, L., del Nozal, M. J., Bernal, J. L., Jiménez, J. J. and Palencia, P. G. (2008). How natural infection by *Nosema ceranae* causes honeybee colony collapse. *Environmental Microbiology* **10**, 2659–2669. doi: 10.1111/j.1462-2920.2008.01687.x.
- Higes, M., Martín-Hernández, R., Garrido-Bailón, E., González-Porto, A. V., García-Palencia, P., Meana, A., Del Nozal, M. J., Mayo, R. and Bernal, J. L. (2009). Honeybee colony collapse due to *Nosema ceranae* in professional apiaries. *Environmental Microbiology Reports* **1**, 110–113. doi: 10.1111/j.1758-2229.2009.00014.x.
- Huang, W., Jiang, J.-H., Chen, Y.-W. and Wang, C.-H. (2007). A *Nosema ceranae* isolate from the honeybee *Apis mellifera*. *Apidologie* **38**, 30–37. doi: 10.1051/apido.
- Huang, W.-F. and Solter, L. F. (2013). Comparative development and tissue tropism in *Nosema apis* and *Nosema ceranae*. *Journal of Invertebrate Pathology* **113**, 35–41. doi: 10.1016/j.jip.2013.01.001.
- Hughes, W. O. H., Eilenberg, J. and Boomsma, J. J. (2002). Trade-offs in group living: transmission and disease resistance in leaf-cutting ants. *Proceedings of the Royal Society B: Biological Sciences* **269**, 1811–1819. doi: 10.1098/rspb.2002.2113.
- Hughes, W. O. H., Petersen, K. S., Ugelvig, L. V., Pedersen, D., Thomsen, L., Poulsen, M. and Boomsma, J. J. (2004). Density-dependence and within-host competition in a semelparous parasite of leaf-cutting ants. *BMC Evolutionary Biology* **4**, 45. doi: 10.1186/1471-2148-4-45.
- Hughes, W. O. H., Bot, A. N. M. and Boomsma, J. J. (2010). Caste-specific expression of genetic variation in the size of antibiotic-producing glands of leaf-cutting ants. *Proceedings Biological Sciences/The Royal Society* **277**, 609–615. doi: 10.1098/rspb.2009.1415.
- Johnson, M. B., Lafferty, K. D., van Oosterhout, C. and Cable, J. (2011). Parasite transmission in social interacting hosts: monogenean epidemics in guppies. *PLoS ONE* **6**, e22634. doi: 10.1371/journal.pone.0022634.
- Klee, J., Besana, A. M., Genersch, E., Gisder, S., Nanetti, A., Tam, D. Q., Chinh, T. X., Puerta, F., Ruz, J. M., Kryger, P., Message, D., Hatjina, F., Korpela, S., Fries, I. and Paxton, R. J. (2007). Widespread dispersal of the microsporidian *Nosema ceranae*, an emergent pathogen of the western honey bee, *Apis mellifera*. *Journal of Invertebrate Pathology* **96**, 1. doi: 10.1016/j.jip.2007.02.014.
- Knell, R. J., Begon, M. and Thompson, D. J. (1996). Transmission dynamics of *Bacillus thuringiensis* infecting *Plodia interpunctella*: a test of the mass action assumption with an insect pathogen. *Proceedings Biological Sciences/The Royal Society* **263**, 75–81. doi: 10.1098/rspb.1996.0013.
- Knell, R. J., Begon, M. and Thompson, D. J. (1998). Transmission of *Plodia interpunctella* granulosis virus does not conform to the mass action

- model. *Journal of Animal Ecology* **67**, 592–599. doi: 10.1046/j.1365-2656.1998.00219.x.
- Lloyd-Smith, J. O., Schreiber, S. J., Kopp, P. E. and Getz, W. M.** (2005). Superspreading and the effect of individual variation on disease emergence. *Nature* **438**, 355–359. doi: 10.1038/nature04153.
- Martin-Hernandez, R., Meana, A., Garcia-Palencia, P., Marin, P., Botias, C., Garrido-Bailon, E., Barrios, L. and Higes, M.** (2009). Effect of temperature on the biotic potential of honeybee microsporidia. *Applied and Environmental Microbiology* **75**, 2554–2557. doi: 10.1128/aem.02908-08.
- Matthews, L., Reeve, R., Woolhouse, M. E. J., Chase-Topping, M., Mellor, D. J., Pearce, M. C., Allison, L. J., Gunn, G. J., Low, J. C. and Reid, S. W. J.** (2009). Exploiting strain diversity to expose transmission heterogeneities and predict the impact of targeting supershedding. *Epidemics* **1**, 221–229. doi: 10.1016/j.epidem.2009.10.002.
- Mayack, C. and Naug, D.** (2009). Energetic stress in the honeybee *Apis mellifera* from *Nosema ceranae* infection. *Journal of Invertebrate Pathology* **100**, 185–188. doi: 10.1016/j.jip.2008.12.001.
- McCallum, H., Barlow, N. and Hone, J.** (2001). How should pathogen transmission be modelled? *Trends in Ecology and Evolution* **16**, 295–300. doi: [http://dx.doi.org/10.1016/S0169-5347\(01\)02144-9](http://dx.doi.org/10.1016/S0169-5347(01)02144-9).
- McCallum, H., Barlow, N. and Hone, J.** (2002). Modelling transmission: mass action and beyond. *Trends in Ecology and Evolution* **17**, 64–65. doi: 10.1016/S0169-5347(01)02399-0.
- Moore, J.** (1995). The behavior of parasitized animals when an ant... is not an ant. *BioScience* **45**, 89–96.
- Naug, D. and Camazine, S.** (2002). The role of colony organization on pathogen transmission in social insects. *Journal of Theoretical Biology* **215**, 427–439. doi: 10.1006/jtbi.2001.2524.
- Naug, D. and Gibbs, A.** (2009). Behavioral changes mediated by hunger in honeybees infected with *Nosema ceranae*. *Apidologie* **40**, 595–599. doi: 10.1051/apido/2009039.
- Naug, D. and Smith, B.** (2007). Experimentally induced change in infectious period affects transmission dynamics in a social group. *Proceedings Biological Sciences/The Royal Society* **274**, 61–65. doi: 10.1098/rspb.2006.3695.
- Nowak, M. A. and May, R. M.** (1994). Superinfection and the evolution of parasite virulence. *Proceedings Biological Sciences/The Royal Society* **255**, 81–89. doi: 10.1098/rspb.1994.0012.
- Nunn, C. L., Lindenfors, P., Pursall, E. R. and Rolff, J.** (2009). On sexual dimorphism in immune function. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **364**, 61–69. doi: 10.1098/rstb.2008.0148.
- O'Donnell, S. and Beshers, S. N.** (2004). The role of male disease susceptibility in the evolution of haplodiploid insect societies. *Proceedings of the Royal Society B: Biological Sciences* **271**, 979–983. doi: 10.1098/rspb.2004.2685.
- Paxton, R. J., Klee, J., Korpela, S. and Fries, I.** (2007). *Nosema ceranae* has infected *Apis mellifera* in Europe since at least 1998 and may be more virulent than *Nosema apis*. *Apidologie* **38**, 558–565. doi: 10.1051/apido.
- Perkins, S. E., Cattadori, I. M., Tagliapietra, V., Rizzoli, A. P. and Hudson, P. J.** (2003). Empirical evidence for key hosts in persistence of a tick-borne disease. *International Journal for Parasitology* **33**, 909–917. doi: 10.1016/S0020-7519(03)00128-0.
- Poulsen, M., Hughes, W. O. H. and Boomsma, J.** (2006). Differential resistance and the importance of antibiotic production in *Acromyrmex echinator* leaf-cutting ant castes towards the entomopathogenic fungus *Aspergillus nomius*. *Insectes Sociaux* **53**, 349.
- Prendini, L., Weygoldt, P. and Wheeler, W. C.** (2005). Systematics of the group of African whip spiders (Chelicerata: Amblypygi): evidence from behaviour, morphology and DNA. *Organisms Diversity and Evolution* **5**, 203–236. doi: 10.1016/j.ode.2004.12.004.
- R Development Core Team** (2012). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Richard, F.-J., Aubert, A. and Grozinger, C. M.** (2008). Modulation of social interactions by immune stimulation in honey bee, *Apis mellifera*, workers. *BMC Biology* **6**, 50. doi: 10.1186/1741-7007-6-50.
- Richard, F.-J., Holt, H. L. and Grozinger, C. M.** (2012). Effects of immunostimulation on social behavior, chemical communication and genome-wide gene expression in honey bee workers (*Apis mellifera*). *BMC Genomics* **13**, 558. doi: 10.1186/1471-2164-13-558.
- Roberts, K. E. and Hughes, W. O. H.** (2014). Immunosenescence and resistance to parasite infection in the honey bee, *Apis mellifera*. *Journal of Invertebrate Pathology*, in press.
- Rolff, J.** (2002). Bateman's principle and immunity. *Proceedings Biological Sciences/The Royal Society* **269**, 867–872. doi: 10.1098/rspb.2002.1959.
- Rosengaus, R. B., Traniello, J. F. A., Lefebvre, M. L. and Carlock, D. M.** (2000). The social transmission of disease between adult male and female reproductives of the dampwood termite *Zootermopsis angusticollis*. *Ethology Ecology and Evolution* **12**, 419–433.
- Rueppell, O., Hayworth, M. K. and Ross, N. P.** (2010). Altruistic self-removal of health-compromised honey bee workers from their hive. *Journal of Evolutionary Biology* **23**, 1538–1546. doi: 10.1111/j.1420-9101.2010.02022.x.
- Ruiz-González, M. X. and Brown, M. J. F.** (2006). Honey bee and bumblebee trypanosomatids: specificity and potential for transmission. *Ecological Entomology* **31**, 616–622. doi: 10.1111/j.1365-2311.2006.00823.x.
- Ryder, J., Miller, M. R., White, A., Knell, R. J. and Boots, M.** (2007). Host-parasite population dynamics under combined frequency- and density-dependent transmission. *Oikos* **116**, 2017–2026. doi: 10.1111/j.2007.0030-1299.15863.x.
- Schmid-Hempel, P.** (1998). *Parasites in Social Insects*. Princeton University Press, USA.
- Shykoff, J. A. and Schmid-Hempel, P.** (1991). Parasites and the advantage of genetic variability within social insect colonies. *Proceedings of the Royal Society B: Biological Sciences* **243**, 55–58. doi: 10.1098/rspb.1991.0009.
- Smart, M. D. and Sheppard, W. S.** (2012). *Nosema ceranae* in age cohorts of the western honey bee (*Apis mellifera*). *Journal of Invertebrate Pathology* **109**, 148–151. doi: 10.1016/j.jip.2011.09.009.
- Smith, M. L.** (2012). The honey bee parasite *Nosema ceranae*: transmissible via food exchange? *PLoS ONE* **7**, e43319. doi: 10.1371/journal.pone.0043319.
- Traniello, J. F. A., Rosengaus, R. B. and Savoie, K.** (2002). The development of immunity in a social insect: evidence for the group facilitation of disease resistance. *Proceedings of the National Academy of Sciences* **99**, 6838–6842. doi: 10.1073/pnas.102176599.
- Traver, B. E. and Fell, R. D.** (2011a). *Nosema ceranae* in drone honey bees (*Apis mellifera*). *Journal of Invertebrate Pathology* **107**, 234–236. doi: 10.1016/j.jip.2011.05.016.
- Traver, B. E. and Fell, R. D.** (2011b). Prevalence and infection intensity of *Nosema* in honey bee (*Apis mellifera* L.) colonies in Virginia. *Journal of Invertebrate Pathology* **107**, 43–49. doi: 10.1016/j.jip.2011.02.003.
- Wilson, K., Knell, R., Boots, M. and Koch-Osborne, J.** (2003). Group living and investment in immune defence: an interspecific analysis. *Journal of Animal Ecology* **72**, 133–143. doi: 10.1046/j.1365-2656.2003.00680.x.
- Woolhouse, M. E. J., Dye, C., Etard, J. F., Smith, T., Charlwood, J. D., Garnett, G. P., Hagan, P., Hii, J. K. L., Ndhlovu, P. D., Quinnell, R. J., Watts, C. H., Chandiwana, S. K. and Anderson, R. M.** (1997). Heterogeneities in the transmission of infectious agents: implications for the design of control programs. *Proceedings of the National Academy of Sciences USA* **94**, 338–342. doi: 10.1073/pnas.94.1.338.